



Evaluation of Some Biomarkers Associated with Anemia in Patients who infected with *Giardia intestinalis*

Adel Hassan Jasim^{1*} , Ikhlās Abbas Marhoon¹

¹Biology Department, College of education, University of Al-Qadisiyah, Al-Diwaniyah-Iraq

Abstract

Introduction: Protozoan infections cause a major part of morbidity in developing countries with poor sanitation and water systems. One of the most prominent protozoan infections is *Giardia intestinalis*, which is a well-known intestinal infectious agent, triggering a host of gastrointestinal problems and malabsorption syndromes, the infection causing structural and functional changes to the intestinal mucosa. These changes to the intestinal mucosa result malabsorption of critical nutrients, such as iron, folate, and the vitamin B12. The study aims to identify the changes in some of hematological and biochemical parameters pertaining to anemia.

Methods: A 200 fecal samples were examine from both sex (100 male and 100 female) in the same ages to reveal a *G. intestinalis* infection. The patients were classified into four groups: *G. intestinalis* infection, Anemia persons, *G. intestinalis* infection and anemia together, and healthy persons as a control; then a blood samples were collected from the forth groups, that attending to parasitological lab. At General health laboratories in Holy Karbala Province – Iraq, from October 2025 until March 2026. A five ml of blood samples were taken and divided into two parts: 2 ml of each sample placed in coagulant tubes and the remaining (3 ml) placed in anticoagulant tubes. The samples were measured and evaluated the mean±SD and LSD test at $P \leq 0.05$ for hematological parameters (RBC, Hb, PCV, MCH, MCHC and WBCs), and some biochemical markers related with Hematopoiesis: Vit. D₃, Vit. B₁₂, Ferritin and serum iron.

Results: The results showed *G. intestinalis* infection effect on hematological parameters and biochemical markers related with hematopoiesis. Anemia in both male and female patients infected with *Giardia* and even some patients who did not present with anemia showed a reduction in RBC, Hb and PCV compared to the control group. Biochemical analysis showed that the infected and anemic group had low values of serum iron, ferritin and vitamin B12 compared to the control group, who had higher levels. Vitamin D3 levels were also showed a lower values in the group with *G. intestinalis* infection plus anemia compared with control group.

Conclusion: The study suggests that infection with *G. intestinalis* causes notable changes in some of hematological and biochemical parameters pertaining to anemia. These changes mirror the relationships between intestinal protozoan infection with blood parameters and nutrient absorption. Even in some patients the combination of *G. intestinalis* infection with anemia presented with the most changes in some parameters. All this changes and the already existing altered blood biochemical and hematological parameters in patients with giardiasis underline the infection's complications.

Keywords: Intestinal protozoa, Anemia, *Giardia intestinalis*, Vit. D₃, Vit. B₁₂

Received: May 13, 2026, Revised: June 11, 2026, Accepted: June 22, 2026, ePublished: July 9, 2026

Introduction

In developing nations, intestinal protozoan infections are a major cause of food and water-borne illnesses. Protozoan infections cause a major part of morbidity in developing countries with poor sanitation and water systems (1). One of the most prominent protozoan infections is *Giardia intestinalis* (*Giardia lamblia*), which is a well-known intestinal infectious agent, triggering a host of gastrointestinal problems and malabsorption syndromes. *Giardia lamblia* infections primarily occur in the small intestine, leading to parasitic attachment to the epithelial layer of the intestine, causing structural and functional changes to the intestinal mucosa (2, 3). These changes to the intestinal mucosa result in the malabsorption of critical nutrients, such as iron, folate, and the vitamin B12 (4, 5). Intestinal protozoan infections cause chronic inflammation of the intestines which contribute to nutrient deficiencies and affect the blood (6).

There have been multiple research reports that appear to suggest the presence of intestinal protozoan parasites

results in noticeable alterations of certain hematological markers that is, hemoglobin levels and mean corpuscular volume as a consequence of their effects on the nutrition and physiology of the infested individual (7). Further, impaired iron absorption and the inflammatory response can ultimately contribute to the development of anemia which is considered a worldwide public health concern across many demographic groups (5, 8).

Holy Karbala province is a region where the aforementioned environmental and demographic elements may aid in the transmission of intestinal protozoan parasites via means of population mobility and problems associated with sanitary systems (9,10).

Therefore, it will be quite relevant to evaluate the various biomarkers associated with anemia in regards to *G. intestinalis* infection within this particular population (2).

This study highlights to effectiveness of *G. intestinalis* infection in some of hematological and biochemical parameters pertaining to anemia, such as the number



of RBCs, Hb, PCV, MCH, MCHC and WBCs, and, and some biochemical markers related with Hematopoiesis Vit. D₃, Vit. B₁₂, Ferritin and serum iron, compared with healthy persons (control group).

Materials and Methods

Study Design and Population

This study aims to assess certain biomarker related to anemia in *G. intestinalis* infected patients in Karbala. The study population consisted of patients visiting health centers and laboratories and was divided into four categories: patients with Giardia, patients with anemia, patients with both Giardia infection and anemia, and a healthy control group. The study included both males and females in the same ages. The grouping of participants was done based on lab identification of Giardia infection and a blood test for anemia.

Sample Collection and Diagnosis

Stool samples were taken from participants and analyzed using standard lab methods for Giardia infections. Identification of Giardia infections was based on sample detection of stages of parasites. All participants provided blood samples for hematological and biochemical analysis. Hematology parameters were analyzed through routine laboratory techniques.

Hematological Parameters

The following hematological parameters were evaluated (7, 9):

1. Red Blood Cell count (RBC)
2. Hemoglobin concentration (Hb)
3. Packed Cell Volume (PCV)
4. Mean Corpuscular Hemoglobin (MCH)
5. Mean Corpuscular Hemoglobin Concentration (MCHC)

White Blood Cell count (WBC) The parameters listed were employed to evaluate the hematological condition of the subjects and to identify the existence and classification of anemia. Biochemical parameters In addition to the hematological indices, the following biochemical parameters related to anemia were also assessed.

Biochemical Parameters

In addition to hematological indices, the following biochemical markers associated with anemia were measured: Serum Iron, Serum Ferritin, Vitamin D3 and Vitamin B12 (6, 7). Theoretical frameworks cite the association of the selected biomarkers with iron metabolism and anemia.

Statistical Analysis

The means and standard deviations are presented as MSD. The statistical evaluations of the study groups were done and the Least Significant Difference (LSD) method was applied to identify the significant differences between the groups. The significance level was set based on the P-values in the statistical tables (11).

Results

Hematological Parameters in Male Patients

Mean $\pm$ standard deviation (MSD), LSD and P values evaluating statistical significance are provided for the four groups: Giardia infected patients, anemic patients, patients with Giardia plus anemia, and the healthy (control group), for comparison of the male participant hematological parameters.

Compared to other groups, the control group had the highest values in RBC, Hb, PCV, MCH, MCHC, and WBC. The control group had the marker showing a significant difference from the Giardia infected and anemic groups. In the Giardia only infected group, the control group had marked b due to significant reductions in RBC, Hb, and PCV. The anemia group had similar reductions in RBC, Hb, and PCV compared to the control group, and the differences were significant in light of the reported LSD and P values.

The group with both Giardia infection and anemia showed further reduction in hematological parameters compared with the control group. The statistical analysis indicated a significant differences among the groups, as demonstrated by the different letter annotations in the Table 1, These values indicate significant differences between the study groups according to the reported P values.

Biochemical Parameters in Male Patients

The biochemical parameters related to anemia were also evaluated among male participants.

These included Vitamin D3, Vitamin B12, Ferritin, and serum Iron.

According to results in Table 2, Vitamin D3, the control group recorded the highest value (42.9), marked with the letter (a), while the Giardia infection plus anemic group showed lower values (13.8) marked with (b), indicating a significant differences among groups.

Vitamin B12 levels in the control group reached 484.7 and were marked with (a), while the other groups showed lower values marked with (b), indicating a statistical significance.

Participants in the control group showed the highest ferritin levels (71.6) and lower values in the infected and anemic groups in (b). The control group also showed a serum iron level of 83.9, and the remaining groups showed lower values marked with (b). The results show a significant changes in both metabolic and hematological biomarkers in male patients infected with Giardia intestinalis and especially when there's anemic co-morbidity.

Hematological Parameters in Female Patients

The hematological parameters of female participants were analyzed and compared among four groups: patients infected with Giardia, patients with anemia, patients with both Giardia infection and anemia, and the healthy control group. The results are expressed as

mean $\pm$ standard deviation (MSD), with LSD and P values used to determine statistical significance.

The control group (Table 3) showed the highest values in all hematological parameters and was marked with the letter (a), indicating a significant differences compared with the other groups. In the group infected with *Giardia* only, the values of RBC, Hb, PCV, MCH, MCHC, and WBC were lower than those of the control group and were marked mainly with the letter (b), indicating statistically significant differences.

The anemia group also showed reduced hematological values compared with the control group. Some parameters were marked with (bc), indicating intermediate values between the infected and control groups but still significantly different according to the reported statistical analysis.

The group with both *Giardia* infection and anemia demonstrated further reductions in several hematological indices compared with the control group, and these were marked with different letters (b or c), confirming the

presence of significant differences among groups.

Biochemical Parameters in Female Patients

In the results which showed in Table 4, there was notable inconsistency for the levels of vitamin D3 among the groups. The control group recorded the highest value (41.5), labeled (a), while the infected and anemic groups showed lower values labeled (b) or (c) showing a significant differences.

For vitamin B12, control group values reached a total of 458.3 and were labeled with an (a). The other groups showed lower values, mostly labeled with a (b) showing a significant difference statistically compared to controls. Ferritin values were highest for the control group (69.8), where the infected and anemic groups had lower values labeled (b) showing significant differences. The levels of serum iron was also different among the groups where the control group had 82.8, while the infected and anemic groups had lower values labeled (b) or (c) showing a significant difference statistically.

Table 1. Effects of *G. intestinalis* infection on Hematological Parameters in Male Patients According to Study Groups

Measured hematological parameters	Study groups (Mean $\pm$ SE)				LSD P $\leq$ 0.05
	G. intestinalis + Anemia	G. intestinalis	Anemia	Control	
Total RBCs ($\times 10^3$ / ml)	3.46 $\pm$ 1.28 B	4.12 $\pm$ 1.82 B	3.41 $\pm$ 1.47 B	5.14 $\pm$ 1.81 a	0.83
Hb (gm / dl)	9.0 $\pm$ 0.9 C	11.5 $\pm$ 1.9 B	10.1 $\pm$ 1.3 B	14.5 $\pm$ 2.1 a	2.3
PCV %	30.2 $\pm$ 1.6 C	38.2 $\pm$ 4.1 B	33.6 $\pm$ 3.6 Bc	45.1 $\pm$ 5.1 a	7.5
MCH ($\times 10^3$ / ml)	20.73 $\pm$ 2.2 B	24.92 $\pm$ 3.8 B	23.68 $\pm$ 3.6 B	32.18 $\pm$ 4.1 a	5.1
MCHC ($\times 10^3$ / ml)	298.9 $\pm$ 10.9 B	322.8 $\pm$ 14.8 B	312.3 $\pm$ 13.4 Bc	355.6 $\pm$ 19.8 A	22.8
WBCs ($\times 10^3$ / ml)	13.68 $\pm$ 2.95 B	12.39 $\pm$ 2.92 B	9.13 $\pm$ 2.1 B	7.45 $\pm$ 1.83 A	1.93

Table 2. Effects of *G. intestinalis* infection on Biochemical Biomarkers in Male Patients According to Study Group

Biochemical Parameters	Study groups (Mean $\pm$ SE)				LSD P $\leq$ 0.05
	G.intestinalis + Anemia	G.intestinalis	Anemia	Control	
Vit. D3	13.8 $\pm$ 3.81 B	22.7 $\pm$ 4.16 B	21.3 $\pm$ 4.23 B	42.9 $\pm$ 5.92 A	9.15
Vit. B12	228.1 $\pm$ 63.7 B	305.4 $\pm$ 85.8 B	320.3 $\pm$ 97.6 B	484.7 $\pm$ 128.7 A	123.4
Ferritin	11.8 $\pm$ 2.7 B	21.3 $\pm$ 4.2 B	12.4 $\pm$ 2.9 B	71.6 $\pm$ 9.3 A	17.6
Iron	39.7 $\pm$ 11.4 B	55.6 $\pm$ 12.7 B	37.1 $\pm$ 9.4 B	83.9 $\pm$ 16.1 A	23.1

Table 3. Effects of *G. intestinalis* infection on Hematological Parameters in Female Patients According to Study Groups

Measured hematological parameters	Study groups (Mean $\pm$ SE)				LSD P $\leq$ 0.05
	G. intestinalis + Anemia	G. intestinalis	Anemia	Control	
Total RBCs ($\times 10^3$ / ml)	3.39 $\pm$ 1.14 B	3.94 $\pm$ 1.56 B	3.32 $\pm$ 1.35 B	4.98 $\pm$ 1.63 A	0.9
Hb (gm / dl)	8.9 $\pm$ 0.8 C	11.1 $\pm$ 1.6 B	9.9 $\pm$ 1.4 BC	13.9 $\pm$ 1.7 A	2.15
PCV %	29.8 $\pm$ 1.4 C	36.1 $\pm$ 3.9 B	33.1 $\pm$ 1.9 BC	43.2 $\pm$ 4.8 A	5.9
MCH ($\times 10^3$ / ml)	20.13 $\pm$ 1.9 B	24.38 $\pm$ 3.6 B	22.36 $\pm$ 3.1 B	31.82 $\pm$ 3.5 A	5.3
MCHC ($\times 10^3$ / ml)	299.2 $\pm$ 11.5 B	320.1 $\pm$ 15.2 B	308.3 $\pm$ 11.7 B	354.2 $\pm$ 19.1 A	24.3
WBCs ($\times 10^3$ / ml)	13.36 $\pm$ 2.82 B	12.23 $\pm$ 3.16 B	8.75 $\pm$ 1.69 A	7.12 $\pm$ 1.92 A	1.78

Table 4. Effects of *G. intestinalis* infection on Biochemical Biomarkers in Female Patients According to Study Group

Biochemical Parameters	Study groups (Mean $\pm$ SE)				LSD $P \leq 0.05$
	<i>G.intestinalis</i> + Anemia	<i>G.intestinalis</i>	Anemia	Control	
Vit. D3	11.8 $\pm$ 3.59 C	20.3 $\pm$ 3.98 B	17.4 $\pm$ 4.23 BC	41.5 $\pm$ 5.83 A	8.22
Vit. B12	219.3 $\pm$ 59.2 B	298.7 $\pm$ 81.6 B	315.2 $\pm$ 91.4 B	458.3 $\pm$ 112.4 A	125.6
Ferritin	10.5 $\pm$ 2.4 B	19.4 $\pm$ 3.9 B	11.3 $\pm$ 2.8 B	69.8 $\pm$ 9.1 A	18.2
Iron	37.6 $\pm$ 10.3 C	59.8 $\pm$ 10.8 B	35.1 $\pm$ 8.3 BC	82.8 $\pm$ 15.3 A	21.5

Compared with the control group, the findings indicate that female patients infected with *G. intestinalis*, especially those with combined anemia, showed changes in the hematological and biochemical markers.

Discussion

This research focused on the blood and biochemical markers associated with anemia due to the infection of *Giardia intestinalis* among individuals in Karbala. The findings indicated a noteworthy distinction across many variables in relation to infected and anemic individuals, those with comorbidities, and the healthy controls. In both genders, there were notable hematological findings indicating a decreased number of red blood cells (RBC), hemoglobin (Hb) levels, and packed cell volume (PCV) in the infected and anemic groups as opposed to the control group (10,12). The lowest levels of these values were seen in those patients who were coinfecting with anemia and the *G. intestinalis*. This leads to the conclusion that the malabsorption due to the intestinal protozoa (i.e., *Giardia*) infection and the nutritional blood parameters impact those individuals (8).

Given that a significant number of malnourished individuals have mucosal damage in the small intestine, the impaired absorption of iron and other essential nutrients, as indicated in the theoretical background, supports the infection of intestinal protozoa as the mechanism to impair the absorption of iron, folate, and vitamin B12, which are essential for the healthy production of red blood cells (4, 13, 14).

Furthermore, the inflammatory response that occurs during infections caused by intestinal protozoa could affect the body's processing of iron and subsequently lead to anemia (15).

Describe an effect of intestinal protozoa on changes of certain hematological parameters, which indicates a negative effect on the nutritional and physiological condition of the subjects. The biochemical data affirm this (12).

When comparing the infected and anemic groups to the control groups, females and males were found to have a lower serum iron and a lower serum ferritin. The finding agrees with the theory of chronic intestinal infection causing a negative iron balance and subsequently causing anemia (4, 15).

Infected groups also had a lower serum vitamin B12 compared to controls. Previous studies have shown that

intestinal infections can impede the absorption of vitamin B12, which is critical in the process of erythropoiesis (7, 16).

Multiple groups showed differing levels of vitamin D3 within the study, revealing lower levels of the vitamin in both the infected and anemic groups compared to the control group (17).

These results could potentially be due to intestinal disorders that affect the absorption of nutrients and disrupt the balance of metabolism (4, 18).

The male-female differences for particular groups could be due to some physiological differences along with variation in the nutritional status and the overall effect of the infection and anemia combined. Yet, in both groups, the control set showed much higher levels of the studied parameters compared to the infected and anemic groups (19).

The study showed that *Giardia intestinalis* infection caused changes in certain parameters of anemia. It also showed that giardiasis and anemia together caused greater changes in certain parameters which showed the importance of early diagnosis and assessment of nutritional and hematological parameters from the bloodstream of the infected individuals (20-26).

Conclusion

The present study suggests that infection with *Giardia intestinalis* in Karbala causes notable changes in some of the hematological and biochemical parameters pertaining to anemia. Anemia in both male and female patients infected with *Giardia* and even some patients who did not present with anemia showed a reduction in RBC, hemoglobin (Hb), packed cell volume (PCV), and red blood cell indices compared to the control group. Biochemical analysis showed that the infected and anemic group had low values of serum iron, ferritin and vitamin B12 compared to the control group, who had higher levels. Variations of the Vitamin D3 levels were also exhibited in the study groups. These changes mirror the intestinal protozoan infection and blood parameters and nutrient absorption as previously described in the study. Even in some patients the combination of *Giardia intestinalis* infection with anemia presented with the most changes in some parameters. This and the already existing altered blood biochemical and hematological parameters in patients with giardiasis underline the infection's complications. Because of this, early infection

control measures should be implemented to lessen the complications of anemia in the affected patients.

Acknowledgments

The study was done in the pathological and parasitological laboratories at General health laboratories in Holy Karbala Province – Iraq.

Authors' Contribution

Conceptualization: Ikhlas Abbas Marhoon, Adel Hassan Jasim

Data Curation: Ikhlas Abbas Marhoon, Adel Hassan Jasim

Formal Analysis: Ikhlas Abbas Marhoon

Investigation: Adel Hassan Jasim

Methodology: Ikhlas Abbas Marhoon, Adel Hassan Jasim

Project Administration: Ikhlas Abbas Marhoon

Software: Ikhlas Abbas Marhoon, Adel Hassan Jasim

Supervision: Ikhlas Abbas Marhoon

Validation: Ikhlas Abbas Marhoon, Adel Hassan Jasim

Visualization: Adel Hassan Jasim

Writing –Original Draft: Adel Hassan Jasim

Writing –Review & Editing: Ikhlas Abbas Marhoon

Competing Interests

The authors declare that they have no known competing financial interests that could have appeared to influence the work documented in the study.

Data Availability Statement

Data are available upon reasonable requesting.

Ethical Approval

The present study was conducted in accordance with the ethical standards of biomedical research involving human participants. Approval for the collection and analysis of fecal and blood samples was obtained from the General Health Laboratories Ethical Committee in Holy Karbala Province, Iraq. All participants were informed of the study objectives, procedures, and potential risks prior to sample collection. Written informed consent was obtained from each participant (or their legal guardian when applicable). Confidentiality of patient data was strictly maintained, and all samples were anonymized to protect participant identity. The study adhered to the principles outlined in the Declaration of Helsinki (2013 revision) regarding medical research ethics. No invasive procedures beyond routine blood sampling were performed, and all laboratory analyses were carried out under biosafety protocols.

Funding

There is No.

References

- World Health Organization (WHO). Antimicrobial Resistance: Global Report on Surveillance. Geneva: WHO; 2014.
- Huang DB, White AC. An updated review on *Cryptosporidium* and *Giardia*. Gastroenterol Clin North Am 2006;35(2):291-314. doi:10.1016/j.gtc.2006.03.006
- Leung AK, Leung AA, Wong AH, Sergi CM, Kam JK. Giardiasis: an overview. Recent Pat Inflamm Allergy Drug Discov 2019;13(2):134-43. doi:10.2174/1872213x13666190618124901
- Abu-Hadal LS, Al-Humaidy NM, Al-Hameedawi DF. Significant micronutrient imbalance associated with *Entamoeba histolytica* and *Giardia lamblia* infections in children. Academia Open 2026;11(1):1-16. doi:10.21070/acopen.11.2026.13115
- Hassanein F, Masoud IM, Awwad ZM, Abdel-Salam H, Salem M, Shehata AI. Microbial bowel infections-induced biochemical and biological abnormalities and their effects on young Egyptian swimmers. Sci Rep 2023;13(1):4597. doi:10.1038/s41598-023-31708-3
- Al-Musawy FA, Hussein MH. Evaluation of interleukin-1 alpha, lipase and vitamin D3 in patients with giardiasis. Big Data Health Care 2022;2(1):1-4.
- Khorsheed BM, Issa AR. Study of intestinal parasites and their effects on some blood parameters among children suffering from diarrhea in Zakho district-Iraq. Egypt Acad J Biol Sci E Med Entomol Parasitol 2025;17(2):83-95. doi:10.21608/eajbse.2025.454598
- Fattahi Bafghi A, Hezardastan E, Fakhraei M, Norouzi M. Impact of *Giardia lamblia* infection on hematological parameters in children from Urmia, Iran. Int J Med Parasitol Epidemiol Sci 2025;6(4):131-4. doi:10.34172/ijmpes.6213
- Al-Sultany AK, Al-Morshidy KA. An epidemiological study of intestinal parasites in children attending the pediatric teaching hospital in the holy city of Karbala, Iraq. Med J Babylon 2023;20(1):95-100. doi:http://dx.doi.org/10.4103/MJBL.MJBL_276_22
- Muttasher EG, Almayali HM, Hassan AJ. Evaluation of hematological and biochemical parameters in patients with intestinal giardiasis in Al Diwaniyah province/Iraq. Microbes Infect Dis 2025.
- Daniel WW, Cross CL. Biostatistics: A Foundation for Analysis in the Health Sciences. John Wiley & Sons; 2018.
- Al-Abodi HR. Estimation of variability in hematological and biochemical parameters of patients with giardiasis. Rev Bionatura 2023;8(1):80. doi:http://dx.doi.org/10.21931/RB/2023.08.01.80
- Taher HM. Levels estimation of iron, zinc and copper in the serum of children infected with giardiasis. Al-Kitab J Pure Sci 2024;8(2):120-4. doi:10.32441/kjps.08.02.p10
- Damania A, Arévalo-Cortés A, Lopez A, Seco-Hidalgo V, Garcia-Ramon D, Lefoulon E, et al. Intestinal microbiota changes due to *Giardia intestinalis* infections in a longitudinal Ecuadorian birth cohort and impact on cobalamin biosynthesis. medRxiv [Preprint]. October 17, 2024. Available from: <https://www.medrxiv.org/content/10.1101/2024.10.15.24315510v1>.
- Humadi AA, Minnat TR. The hematological and histopathological changes of *Giardia duodenalis* on the intestine in Diyala government. Diyala J Vet Sci 2023;1(1):66-84. doi:10.71375/djvs.2023.01107
- Arbabi M, Esmaili N, Parastouei K, Hooshyar S, Rasti S. Levels of zinc, copper, magnesium elements, and vitamin B12, in sera of school children with giardiasis and enterobiasis in Kashan, Iran. Zahedan J Res Med Sci 2015; 17(11):e3659. doi:10.17795/zjrms-3659
- Osman H, Hassan M, Ahmed M, Ahmed N. Vitamin D estimation in *Giardia lamblia* infected patients in upper Egypt. J Egypt Soc Parasitol 2023;53(1):71-4. doi:10.21608/jesp.2023.297347
- Jaffar ZA, Merdaw MA. Incidence of intestinal parasitic infections among random samples at Al-Aziziyah hospital in Wasit province/Iraq. Baghdad Sci J 2021;18(2):3. doi:10.21123/bsj.2021.18.2.0217
- Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. J Clin Invest 2008;118(4):1311-21. doi:10.1172/jci34261
- Naser SK. Prevalence of *Giardia* infection: a cross-sectional study in Basra province, Iraq. AUIQ Complement Biol Syst 2025;2(3):65-70. doi:10.70176/3007-973X.1047
- Bakr M, Mohammad S, Kadir MA. Distribution of *Giardia lamblia* among local and displaced children in Kirkuk city. Tikrit J Pure Sci 2023;23(9):28-31. doi:10.25130/tjps.v23i9.809
- Garedaghi Y. A review of the importance of stool test in the diagnosis of intestinal parasites. Int J Med Parasitol Epidemiol Sci 2020;1(2):21-4. doi:10.34172/ijmpes.2020.09
- Garedaghi Y, Hassanzadeh Khanmiri H. Prevalence of human intestinal parasites in Tabriz city of Iran during 2018

- and the importance of these parasites in public health. Int J Med Parasitol Epidemiol Sci 2020;1(4):87-90. doi:[10.34172/ijmpes.2020.26](https://doi.org/10.34172/ijmpes.2020.26)
24. Jannati R, Tavakoli Pasand S, Garedaghi Y. Evaluation of different techniques in laboratory diagnosis of intestinal amoebiasis. Int J Med Parasitol Epidemiol Sci 2024;5(1):16-23. doi:[10.34172/ijmpes.3129](https://doi.org/10.34172/ijmpes.3129)
 25. Davoudi Y, Garedaghi Y, Safarmashaei S. Epidemiological study of giardiasis in diarrheic calves in East-Azerbaijan province, Iran. J Anim Vet Adv 2011;10(19):2508-10. doi:[10.3923/javaa.2011.2508.2510](https://doi.org/10.3923/javaa.2011.2508.2510)
 26. Hariri D, Garedaghi Y. Comparison of therapeutic effects of hydroalcoholic extract of asafoetida with metronidazole in mice infected with *Giardia lamblia*. J Zoonotic Dis 2024;8(1):452-9. doi:[10.22034/jzd.2024.17396](https://doi.org/10.22034/jzd.2024.17396)

© 2026 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.